

Planck Scale Effect in the Entropic Force Law

Subir Ghosh

Physics and Applied Mathematics Unit,
Indian Statistical Institute,
203 B.T.Road, Kolkata 700108, India.

Abstract

In this note we generalize the quantum uncertainty relation proposed by Vancea and Santos [7] in the entropic force law, by introducing Planck scale modifications. The latter is induced by the Generalized Uncertainty Principle. We show that the proposed uncertainty relation of [7], involving the entropic force and the square of particle position, gets modified from the consideration of a minimum measurable length, (which can be the Planck length).

Very recently Verlinde [1] has conjectured that the origin of Newtonian gravity and (second) law of dynamics might be entropic in nature. This means that the conventional forces can originate from maximization of entropy principle as in thermodynamics of macroscopic systems. This framework attempts to establish thermodynamics as the fundamental principle. These ideas are further strengthened by the deep connection between thermodynamics and black hole physics, as advocated by previous workers [2, 3, 4, 5]. A number of subsequent developments [6] in various directions have been reported after Verlinde's work.

Deep physical insight coupled with very simple algebra has led [1] from the entropy principle to the second law of Newtonian dynamics,

$$P = ma \tag{1}$$

where the force P on a particle is related to its mass m and acceleration a . This is derived from the the first law of thermodynamics,

$$\Delta W = T \Delta S = F \Delta x \quad (2)$$

where the variation in energy W of a macroscopic system is expressed in terms of its equilibrium temperature T and change in entropy S . Furthermore, as is customary in thermodynamics, this can also be identified as the work done by a generalized force F for the displacement Δx . F is termed as the entropic force. The essential cog in this analysis is the conjecture that there is a variation in entropy associated with the holographic screen, (that separates the emerged spacetime from the sector to be emerged), as a particle of mass m approaches very close (of the order of the particle Compton length $l_c = \hbar/mc$) to the screen. Explicitly it is postulated that [1]

$$\Delta S = 2\pi k_B \frac{\Delta x}{l_c}, \quad (3)$$

where Δx is the distance of the particle from the screen. Furthermore, Verlinde [1] identifies the thermodynamic temperature T in (2) to the Unruh temperature [3] T_U

$$T_U = \frac{\hbar a}{2\pi k_B c}, \quad (4)$$

where a denotes the acceleration of the observer who experiences T_U . Combining all these a simple algebra leads to Newton's law (1).

In a recent paper Vancea and Santos [7] have opened another line of thought. They have pointed out [7] that the postulate (3) as well as the expression for Unruh temperature (4) are essentially quantum in nature, with the explicit presence of $\hbar$, although $\hbar$ does not show up in the final outcome, Newton's classical law of motion. Indeed, as far as the latter is concerned, this is as it should be but one can expect quantum corrections to Newton's second law of motion and the law of gravitation. It has been emphasized by Padmanabhan already [5] that gravity is intrinsically quantum in the holographic approach as the "classical limit" $\hbar \rightarrow 0$ leads to a divergence in the Newton constant. However, more importantly

for our purpose, is the introduction of Planck length L_P in [5] as a fundamental scale for counting the number of micro cells in the holographic screen that yields the entropy. In [7] the possibility of quantum correction to Newton's law of motion (and gravitation) has been speculated. In the present work we have generalized the work in [7] by incorporating the other essential length scale L_P in the generalized Newton's law.

More explicitly, in [7] the authors have considered the possibility of having an uncertainty in the entropy in the holographic screen originating from the fact that there are inherent *quantum* uncertainties in the position and momentum of a particle. This comes primarily because the information (or entropy) associated with the screen depends linearly on the distance of the test particle from the screen. Keeping this in mind, in [7], the relation (3) has been generalized to

$$\delta S = 2\pi k_B \left(\frac{\delta x}{l_c} + \frac{\delta p}{mc} \right). \quad (5)$$

In fact another alternate form of δS has also been suggested in [7] but we choose here the minimal form. One can think of the denominators as $l_c = \hbar/(mc)$ (for δx) and $\hbar/l_c = mc$ (for δp). In [7] the variations δx and δp are to be considered as quantum uncertainties, obeying the Heisenberg uncertainty relation

$$\delta x \delta p \geq \frac{\hbar}{2}. \quad (6)$$

Indeed in the classical case δx reduces to the separation Δx between the screen and the particle and $\delta p = 0$ as the Heisenberg uncertainty relation does not apply. Then (5) reduces to the Verlinde formula (3). Otherwise, one can replace δp in (5) by $\delta p = \hbar/(2\delta x)$ to obtain a quantum corrected Newton's second law of motion [7]

$$F(\delta) = ma + \frac{\hbar}{2m} \left(\frac{\hbar a}{c^2} - p \right) (\delta x)^{-2}. \quad (7)$$

The above yields an uncertainty relation, first proposed in [7],

$$\delta F (\delta x)^2 \geq \frac{\hbar}{2m} \left(\frac{\hbar a}{c^2} - p \right). \quad (8)$$

Furthermore, using the classical (Newtonian) expression for gravitational acceleration $a = (GM)/R^2$ a quantum corrected Newton's law for gravitation is also obtained [7], to first

order in $\hbar$,

$$F = G \frac{Mm}{R^2} + \frac{\hbar}{2m} \left(G \frac{\hbar M}{R^2 c^2} - p \right) (\delta x)^{-2}. \quad (9)$$

In this perspective, once the quantum correction has been introduced, it is indeed natural to consider the Generalized Uncertainty Principle (GUP) [8] instead of the Heisenberg relation (6). In recent years there has been a lot of activity, studying the consequences of GUP (in non-relativistic [9] and cosmological [10] contexts) since it introduces a minimum length scale, generally considered to be the Planck length $l_P^2 = G\hbar/c^3$. Implications of GUP in the context of Information Theory and Holography, that are very relevant for the present work, was studied in [11]. A non-trivial prediction of generic quantum gravity theories (such as String Theory) and black hole physics is the existence of a minimum measurable length [8]. Heuristically, a minimum length cut off is needed to avoid the paradox that localization of an event below Planck length can generate sufficient energy density to create a black hole, thus rendering the event itself unobservable. For the more numerically minded, the Schwarzschild radius near the Planck scale, $l_s \sim (M_P G)/c^2 \sim \sqrt{(G\hbar)/c^3}$ becomes comparable to the Compton wavelength, $l_c \sim \hbar/(M_P c) \sim \sqrt{(G\hbar)/c^3}$. Incidentally both are of the order of the Planck length. This cherished length scale can be induced by the GUP, of the following form first provided in [12],

$$\delta x_i \delta p_i \geq \frac{\hbar}{2} [1 + \beta((\delta p)^2 + \langle p \rangle^2) + 2\beta(\delta p_j^2 + \langle p_j \rangle^2)] , \quad i = 1, 2, 3 \quad (10)$$

where $p^2 = \sum_{j=1}^3 p_j p_j$, $\beta \sim 1/(M_P c)^2 = \ell_P^2/2\hbar^2$, M_P = Planck mass, and $M_P c^2$ = Planck energy $\approx 1.2 \times 10^{19}$ GeV. Subsequently, the Hilbert space representation of the generalized uncertainty relation was developed in [13]. This is indeed crucial for formulating the quantum mechanics in a phase space with a non-commutative structure that is compatible with the above GUP.

Our main result, in this short note, is to show that the minimum length scale, via the GUP, modifies the new force-position uncertainty relation (8), suggested in [7]. In particular

we find,

$$\delta F(\delta x)^2 \geq \nu \frac{\hbar}{2m} \left(\frac{\hbar a}{c^2} - p \right), \quad (11)$$

where $\nu = 1 + \frac{\hbar^2 \beta}{4\delta x^2}$ is the GUP induced β -dependent correction. Taking the minimum value of δx to be l_P we find the maximum value of ν to be $\nu \sim 1 + 5/4$. (The exact numerical factor should not be taken too seriously.) We will derive this relation in rest of the note.

We start by simplifying the GUP to one dimension,

$$\delta x \delta p \geq \frac{\hbar}{2} (1 + \beta(\delta p)^2). \quad (12)$$

Solving the saturation condition to $O(\beta)$ we find

$$\delta p = \frac{\hbar}{2\delta x} \left(1 + \frac{\hbar^2 \beta}{4} \right). \quad (13)$$

and demanding the reality of δp yields the inequality,

$$\delta x \geq \sqrt{\beta} \hbar, \quad (14)$$

with $\delta x = \sqrt{\beta} \hbar$ being the minimum measurable length. This is taken as Planck length L_P . Thus, following the same steps as in [7] we obtain the expression for entropic force consistent with GUP (12),

$$F = ma + \frac{\hbar}{2mc^2(\delta x)^2} (a\hbar - pc^2) \left(1 + \frac{\hbar^2 \beta}{4(\delta x)^2} \right). \quad (15)$$

As before [7], defining the uncertainty in F to be $\delta F = F - ma$ [7] we obtain

$$\delta F(\delta x)^2 \geq \frac{\hbar}{2mc^2} (a\hbar - pc^2) \left(1 + \frac{\hbar^2 \beta}{4(\delta x)^2} \right) \equiv \nu \frac{\hbar}{2mc^2} (a\hbar - pc^2), \quad (16)$$

in conjunction with the minimum measurable length $\delta x \geq \sqrt{\beta} \hbar \sim L_P$. This is the major result of our paper. The effect of the β -correction can be seen from the relation,

$$(\delta x)_\beta^2 = \frac{1}{2} [(\delta x^2) + (\delta x)^2 \sqrt{1 + \frac{\hbar^2 \beta}{(\delta x)^2}}], \quad (17)$$

where we considered the saturation values of δx^2 for the two cases, ($(\delta x)^2$ for $\beta = 0$ [7] and $(\delta x)_\beta^2$ for $\beta \neq 0$), for the same δF .

Incidentally this ties up nicely with the idea of Padmanabhan [5] that a naive $\hbar \rightarrow 0$ limit is not acceptable as it leads to a diverging Newton's constant. On the other hand the new uncertainty relation (8), proposed in [7], has the possibility of generating large uncertainty $\delta F \rightarrow \infty$ as $\delta x \rightarrow 0$. However, our analysis shows that the minimum length scale L_P once comes again to the rescue by weakening the uncertainty relation of [7] thus allowing finite values of δF .

Let us conclude by pointing out the significance of our analysis in the context of experimental/observational validity to ascertain the consistency of the entropic scenario, its quantum correction, as well as the hypothesis of minimum length scale. For this, two features of our results should be worth mentioning: (i) The correction factor ν in (16) is not damped by the Planck scale. (ii) The purported quantum corrections appear in the classical laws of motion. Hence it is conceivable that one might conduct suitable experiments where additive physical quantities such as mass, energy etc. appear as observables and their cumulative effects might leave a substantial signature of quantum or even the Planckian regime in a classical scenario.

Acknowledgment: It is a pleasure to thank Professor I. V. Vancea for correspondence. I am also grateful to the Referee for comments.

References

- [1] E. P. Verlinde, “On the Origin of Gravity and the Laws of Newton,” arXiv:1001.0785 [hep-th].
- [2] J. D. Bekenstein, “Black holes and entropy,” Phys. Rev. D **7**, 2333 (1973).
- [3] W. G. Unruh, “Notes on black hole evaporation,” Phys. Rev. D **14**, 870 (1976).
- [4] T. Jacobson, “Thermodynamics of space-time: The Einstein equation of state,” Phys. Rev. Lett. **75**, 1260 (1995) [arXiv:gr-qc/9504004].
- [5] T. Padmanabhan, arXiv:0912.3165v2 [gr-qc]
- [6] R. G. Cai, L. M. Cao and N. Ohta, “Friedmann Equations from Entropic Force,” arXiv:1001.3470 [hep-th]; S. W. Wei, Y. X. Liu and Y. Q. Wang, “Friedmann equation of FRW universe in deformed Horava-Lifshitz gravity from entropic force,” arXiv:1001.5238 [hep-th]; Y. Ling and J. P. Wu, “A note on entropic force and brane cosmology,” arXiv:1001.5324 [hep-th]; Y. S. Myung, “Entropic force in the presence of black hole,” arXiv:1002.0871 [hep-th]; Y. X. Liu, Y. Q. Wang and S. W. Wei, “Temperature and Energy of 4-dimensional Black Holes from Entropic Force,” arXiv:1002.1062 [hep-th]; C. Gao, “Modified Entropic Force,” arXiv:1001.4585 [hep-th]; M. Li and Y. Wang, “Quantum UV/IR Relations and Holographic Dark Energy from Entropic Force,” arXiv:1001.4466 [hep-th]; T. Wang, “The Coulomb Force as an Entropic Force,” arXiv:1001.4965 [hep-th]; A. Pesci, “Gravity from the entropy of light,” arXiv:1002.1257 [gr-qc]; R. G. Cai, L. M. Cao and N. Ohta, “Notes on Entropy Force in General Spherically Symmetric Spacetimes,” arXiv:1002.1136 [hep-th]; J. Kowalski-Glikman, “A note on gravity, entropy, and BF topological field theory,” arXiv:1002.1035 [hep-th]; J. W. Lee, H. C. Kim and J. Lee, “Gravity from Quantum Information,” arXiv:1001.5445 [hep-th]; Y. Wang, “Towards a Holographic Description of Inflation and Generation of Fluctuations from Thermodynamics,” arXiv:1001.4786 [hep-th]; Yue Zhao. Entropic

force and its fluctuation from gauge/gravity duality, arXiv:1002.4039 [hep-th], R.A. Konoplya, Entropic force, holography and thermodynamics for static space-times. arXiv:1002.2818 [hep-th] , Francesco Caravelli, Leonardo Modesto, Holographic actions from black hole entropy, arXiv:1001.4364 [gr-qc] , T. Padmanabhan, arXiv:1003.5665 Surface Density of Spacetime Degrees of Freedom from Equipartition Law in theories of Gravity.

- [7] I. V. Vancea, M. A. Santos, arXiv:1002.2454v1 [hep-th].
- [8] D. Amati, M. Ciafaloni, G. Veneziano, Phys. Lett. **B 216** (1989) 41; D.J. Gross, P.F. Mende, Phys. Lett. B 197 (1987) 129; Nucl.Phys. B 303 (1988) 407. K. Konishi, G. Paffuti, P. Provero, Phys.Lett. B 234 (1990) 276; M. Maggiore, Phys. Lett. B **304** (1993) 65 [arXiv:hep-th/9301067]; L. J. Garay, Int. J. Mod. Phys. A **10** (1995) 145 [arXiv:gr-qc/9403008].
- [9] Fabio Scardigli, Phys.Lett. B452 (1999) 39-44 [arXiv:hep-th/9904025], arXiv:gr-qc/0607010 ; arXiv:hep-th/0307174; F. Brau, J. Phys. **A 32** (1999) 7691 [arXiv:quant-ph/9905033]; S. Das and E.C. Vagenas, Phys.Rev.Lett. 101, 221301 (2008); B. Bolen and M. Cavaglià, Gen.Rel.Grav. 37, 1255 (2005).
- [10] Mu-in Park, Phys.Lett.B659:698-702,2008 (arXiv:0709.2307v4 [hep-th]); Tao Zhu, Ji-Rong Ren, Ming-Fan Li, Phys.Lett.B674:204-209,2009 (arXiv:0811.0212v3 [hep-th]).
- [11] A. Kempf, Phys. Rev. Lett. 92, 221301 (2004).
- [12] A. Kempf, J. Phys. A: Math. Gen. 30 2093 (1997).
- [13] A. Kempf, G. Mangano, R. B. Mann, Phys. Rev. **D52** (1995) 1108 [arXiv:hep-th/9412167]; A. Kempf, J.Phys. **A 30** (1997) 2093 [arXiv:hep-th/9604045].